arXiv:math/0401228v2 [math.NT] 6 Mar 2005

Accepted by Proc. Amer. Math. Soc. on March 6, 2005.

BINOMIAL COEFFICIENTS AND QUADRATIC FIELDS

Zhi-Wei Sun

Department of Mathematics, Nanjing University
Nanjing 210093, People's Republic of China
zwsun@nju.edu.cn `http://pweb.nju.edu.cn/zwsun`

Abstract. Let E be a real quadratic field with discriminant $d \not\equiv 0 \pmod p$ where p is an odd prime. For $\rho = \pm 1$ we determine $\prod_{0<c<d,\ (\frac{d}{c})=\rho} \binom{p-1}{\lfloor pc/d \rfloor}$ modulo p^2 in terms of Lucas numbers, the fundamental unit and the class number of E.

1. Introduction

Let p be an odd prime not dividing a positive integer m. A. Granville [G, (1.15)] discovered the remarkable congruence

$$\prod_{0<k<m} \binom{p-1}{\lfloor pk/m \rfloor} \equiv (-1)^{(m-1)(p-1)/2}(m^p - m + 1) \pmod{p^2},$$

where we use $\lfloor x \rfloor$ to denote the integral part of a real number x. Subsequently the present author [S1] determined further $\prod_{0<k<m/2} \binom{p-1}{\lfloor pk/m \rfloor}$ mod p^2. In this paper a more sophisticated result connected with real quadratic fields will be established.

For $A, B \in \mathbb{Z}$ the Lucas sequences $u_n = u_n(A, B)$ and $v_n = v_n(A, B)$ $(n = 0, 1, 2, \ldots)$ are given by

$$\begin{aligned} &u_0 = 0,\ u_1 = 1, \text{ and } u_{n+1} = Au_n - Bu_{n-1} \text{ for } n = 1, 2, 3, \ldots, \\ &v_0 = 2,\ v_1 = A, \text{ and } v_{n+1} = Av_n - Bv_{n-1} \text{ for } n = 1, 2, 3, \ldots. \end{aligned}$$

It is well known that

$$(\alpha - \beta)u_n = \alpha^n - \beta^n \quad \text{and} \quad v_n = \alpha^n + \beta^n \quad \text{for every } n = 0, 1, 2, \ldots,$$

where α and β are the two roots of the equation $x^2 - Ax + B = 0$. Also, for any odd prime p we have $u_p \equiv (\frac{\Delta}{p}) \pmod p$ and $v_p \equiv A \pmod p$, where $\Delta = A^2 - 4B$ and $(\frac{\cdot}{p})$ denotes the Legendre symbol. (See, e.g., [R, pp. 41-55].) If p is an odd prime not dividing B, then $p \mid u_{p-(\frac{\Delta}{p})}$ since $Au_p + v_p = 2u_{p+1}$ and $Au_p - v_p = 2Bu_{p-1}$.

2000 *Mathematics Subject Classifications*. Primary 11B65; Secondary 11B37, 11B68, 11R11.

The author was supported by the National Science Fund for Distinguished Young Scholars (No. 10425103) and the Key Program of NSF (No. 10331020) in China.

Throughout this paper, for an assertion P we set

$$[P]=\begin{cases}1 & \text{if } P \text{ holds,}\\ 0 & \text{otherwise.}\end{cases} \tag{1.1}$$

Our main result is as follows.

Theorem 1.1. *Let E be a quadratic field with discriminant $d=2^{\alpha}p_1\cdots p_r$ where $\alpha\in\{0,2,3\}$ and $p_1,\ldots,p_r$ are distinct odd primes. Let $\varepsilon=(a+b\sqrt{d})/2$ be the fundamental unit of the field E where $a,b\in\mathbb{Z}$, and $N(\varepsilon)$ be the norm $(a^2-b^2d)/4$ of ε with respect to the field extension $E/\mathbb{Q}$. Let h be the class number of the field E, and p be an odd prime not dividing d. Then, for $\rho=\pm1$ we have*

$$\begin{aligned}\prod_{\substack{0<c<d\\ (\frac{d}{c})=\rho}}\binom{p-1}{\lfloor pc/d\rfloor}\equiv&1+\frac{\varphi(d)}{2}\bigg((\alpha+[\alpha>0])(2^{p-1}-1)+\sum_{0<i\leqslant r}\frac{p_i^p-p_i}{p_i-1}\bigg)\\ &+\frac{\rho}{2}\left(\frac{d}{p}\right)^{[N(\varepsilon)=1]}u_{p-(\frac{d}{p})}(a,N(\varepsilon))bdh \pmod{p^2},\end{aligned} \tag{1.2}$$

where φ is Euler's totient function and $(\frac{d}{\cdot})$ is the Kronecker symbol.

Remark. Under the conditions of Theorem 1.1, $d\equiv1 \pmod 4$ if $\alpha=0$, and $d/4\equiv3 \pmod 4$ if $\alpha=2$; also p divides $bu_{p-(\frac{d}{p})}(a,N(\varepsilon))$ since for $p\nmid b$ we have

$$\left(\frac{a^2-4N(\varepsilon)}{p}\right)=\left(\frac{b^2d}{p}\right)=\left(\frac{d}{p}\right).$$

Example. Each of the quadratic fields $\mathbb{Q}(\sqrt{13}),\mathbb{Q}(\sqrt{21}),\mathbb{Q}(\sqrt{6}),\mathbb{Q}(\sqrt{7})$ has class number 1, and their fundamental units are

$$\frac{3+\sqrt{13}}{2},\ \frac{5+\sqrt{21}}{2},\ 5+2\sqrt{6}=\frac{10+2\sqrt{24}}{2},\ 8+3\sqrt{7}=\frac{16+3\sqrt{28}}{2}$$

with norms $-1,1,1,1$ respectively; see, e.g., [C, p. 271]. Let p be an odd prime and $\rho\in\{1,-1\}$. If p does not divide 13, 21, 6, and 7, respectively, then Theorem 1.1 gives the congruences

$$\prod_{\substack{0<c<13\\ (\frac{13}{c})=\rho}}\binom{p-1}{\lfloor pc/13\rfloor}\equiv1+\frac{13^p-13}{2}+\rho\frac{13}{2}u_{p-(\frac{13}{p})}(3,-1),$$

$$\prod_{\substack{0<c<21\\ (\frac{21}{c})=\rho}}\binom{p-1}{\lfloor pc/21\rfloor}\equiv1+3(3^p-3)+7^p-7+\rho\left(\frac{21}{p}\right)\frac{21}{2}u_{p-(\frac{21}{p})}(5,1),$$

$$\prod_{\substack{0<c<24\\ 2\nmid c,\ (\frac{6}{c})=\rho}}\binom{p-1}{\lfloor pc/24\rfloor}\equiv1+8(2^p-2)+2(3^p-3)+\rho\left(\frac{6}{p}\right)24u_{p-(\frac{6}{p})}(10,1),$$

$$\prod_{\substack{0<c<28\\ 2\nmid c,\ (\frac{7}{c})=\rho}}\binom{p-1}{\lfloor pc/28\rfloor}\equiv1+9(2^p-2)+7^p-7+\rho\left(\frac{7}{p}\right)42u_{p-(\frac{7}{p})}(16,1)$$

modulo p^2 respectively, where $(\frac{6}{c})$ and $(\frac{7}{c})$ are Jacobi symbols.

We deduce Theorem 1.1 by combining the following two theorems.

Theorem 1.2. *Let $m > 2$ be an integer with the factorization $p_1^{\alpha_1}\cdots p_r^{\alpha_r}$ where $p_1,\ldots,p_r$ are distinct primes and $\alpha_1,\ldots,\alpha_r$ are positive integers. Let p be an odd prime not dividing m. Then*

$$\begin{aligned}&(-1)^{\frac{\varphi(m)}{2}\cdot\frac{p-1}{2}}\left(\frac{p_1}{p}\right)^{[r=1]}\prod_{\substack{0<k<m/2\\(k,m)=1}}\binom{p-1}{\lfloor pk/m\rfloor}\\ \equiv&1+\frac{\varphi(m)}{2}\sum_{i=1}^{r}(\alpha_i p_i-\alpha_i+1)\frac{p_i^{p-1}-1}{p_i-1}\pmod{p^2}.\end{aligned}\tag{1.3}$$

In the next theorem we use the Bernoulli polynomial $B_n(x)$ of degree n and the nth Bernoulli number $B_n = B_n(0)$. Also, we let $\mathbb{P}$ denote the set of all (positive) primes.

Theorem 1.3. *Let E be a real quadratic field with discriminant d and class number h. Let $\varepsilon = (a+b\sqrt{d})/2 > 1$ be the fundamental unit of E where $a,b\in\mathbb{Z}$, and $N(\varepsilon)$ be the norm $(a^2-b^2d)/4$ of ε. Let p be an odd prime not dividing d, and let u stand for $bu_{p-(\frac{d}{p})}(a,N(\varepsilon))$. Then*

$$\sum_{c=1}^{d-1}\left(\frac{d}{c}\right)\left(B_{p-1}\left(\frac{c}{d}\right)-B_{p-1}\right)\equiv\left(\frac{d}{p}\right)^{[N(\varepsilon)=-1]}dh\frac{u}{p}\pmod{p},\tag{1.4}$$

and

$$\prod_{\substack{0<c<d/2\\(c,d)=1}}\binom{p-1}{\lfloor pc/d\rfloor}^{(\frac{d}{c})}\equiv\begin{cases}(\frac{d}{p})(1+\frac{dhu}{2})\pmod{p^2} & \text{if } d=8 \text{ or } d\in\mathbb{P},\\ 1+(\frac{d}{p})^{[N(\varepsilon)=1]}\frac{dhu}{2}\pmod{p^2} & \text{otherwise.}\end{cases}\tag{1.5}$$

Remark. In the case where $d\equiv 1 \pmod 4$ is a prime, (1.4) was proved in [GS] by means of p-adic logarithms and Dirichlet's class number formula (see, e.g., [W]).

In the spirit of R. Crandall and C. Pomerance [CP], Theorems 1.1–1.3 might be of computational interest.

We shall make some preparations in the next section and give proofs of Theorems 1.1–1.3 in Section 3.

2. On the sum $\sum_{\substack{0<k<p\\ m\mid k-r}}\frac{1}{k}$ modulo p

Bernoulli polynomials play important roles in many aspects. The reader is referred to [IR, pp. 228-248] for basic properties, and to [DSS] for a bibliography of related papers.

In this section we prove the following basic result and derive some consequences.

Theorem 2.1. *Let m be a positive integer not divisible by an odd prime p. Then for any $r\in\mathbb{Z}$ we have*

$$\sum_{\substack{k=1\\ k\equiv r\ (\mathrm{mod}\ m)}}^{p-1}\frac{1}{k}\equiv\frac{1}{m}\left(B_{p-1}\left(\left\{\frac{r}{m}\right\}\right)-B_{p-1}\left(\left\{\frac{r-p}{m}\right\}\right)\right)\ (\mathrm{mod}\ p),\tag{2.1}$$

where $\{x\}$ stands for the fractional part of a real number x.

Proof. Applying Lemma 3.1 of [S3] with $k=p-2$, we find that

$$-m\sum_{\substack{j=1\\ j\equiv r\ (\mathrm{mod}\ m)}}^{p-1}\frac{1}{j}\equiv B_{p-1}\left(\frac{p}{m}+\left\{\frac{r-p}{m}\right\}\right)-B_{p-1}\left(\left\{\frac{r}{m}\right\}\right)\ (\mathrm{mod}\ p).$$

For $t=\{(r-p)/m\}$, we have

$$B_{p-1}\left(\frac{p}{m}+t\right)-B_{p-1}(t)=\sum_{l=1}^{p-1}\binom{p-1}{l}B_{p-1-l}\left(\left(\frac{p}{m}+t\right)^l-t^l\right)\equiv 0\ (\mathrm{mod}\ p).$$

(Recall that $B_1=-1/2$ and $B_{2n+1}=0$ for $n=1,2,\ldots$. Also, p divides no denominators of $B_0,B_2,\ldots,B_{p-3}$ by the theorem of Clausen and von Staudt (cf. [IR, pp. 233-236]).) Therefore (2.1) follows. □

Remark. The author first discovered Theorem 2.1 in Sept. 1991 by using Fourier series, and Lemma 3.1 of [S3] was originally motivated by this result.

Corollary 2.1. *Let m and n be positive integers, and let p be an odd prime not dividing m. Then*

$$B_{p-1}\left(\left\{\frac{pn}{m}\right\}\right)-B_{p-1}\equiv m\sum_{r=1}^{n}K_p(r,m)\equiv-\sum_{\substack{k=1\\ p\nmid k}}^{\lfloor pn/m\rfloor}\frac{1}{k}\ \ (\mathrm{mod}\ p),\tag{2.2}$$

where

$$K_p(r,m):=\sum_{\substack{k=1\\ m\mid k-rp}}^{p-1}\frac{1}{k}=\sum_{\substack{l=1\\ m\mid l-(1-r)p}}^{p-1}\frac{1}{p-l}\equiv-K_p(1-r,m)\ (\mathrm{mod}\ p).\tag{2.3}$$

Proof. In view of Theorem 2.1,

$$\begin{aligned} m\sum_{r=1}^{n}K_p(r,m)\equiv&\sum_{r=1}^{n}\left(B_{p-1}\left(\left\{\frac{rp}{m}\right\}\right)-B_{p-1}\left(\left\{\frac{(r-1)p}{m}\right\}\right)\right)\\ \equiv&B_{p-1}\left(\left\{\frac{pn}{m}\right\}\right)-B_{p-1}\pmod p. \end{aligned}$$

On the other hand,

$$-\sum_{r=1}^{n}K_p(r,m)\equiv\sum_{r=1}^{n}\sum_{\substack{k=1\\ m\mid rp-k}}^{p-1}\frac{1}{rp-k}=\sum_{\substack{j=1\\ p\nmid j,\ m\mid j}}^{pn-1}\frac{1}{j}=\sum_{\substack{k=1\\ p\nmid k}}^{\lfloor pn/m\rfloor}\frac{1}{km}\pmod p.$$

So we have (2.2). □

Let p be an odd prime and r be any integer. An explicit congruence for $K_p(r,12)$ mod p appeared in Corollary 3.3 of [S2]. By Theorem 2.1 and [GS, (4)] we can also determine

$$K_p(3+6r,24),\ K_p(5,40),\ K_p(25,40),\ K_p(6,60),\ K_p(36,60)$$

modulo p in terms of some second-order linear recurrences.

For a prime p and any $a\in\mathbb{Z}$ not divisible by p, the Fermat quotient $q_p(a)$ is defined as the integer $(a^{p-1}-1)/p$.

Corollary 2.2. *Let p be an odd prime and let m be a positive integer not divisible by p. Then we have*

$$\sum_{r=1}^{m}rK_p(r,m)\equiv-q_p(m)\pmod p. \tag{2.4}$$

Proof. By Corollary 2.1,

$$\begin{aligned} \sum_{n=1}^{m}\sum_{r=1}^{n}K_p(r,m)\equiv&\frac{1}{m}\sum_{n=1}^{m}\left(B_{p-1}\left(\left\{\frac{pn}{m}\right\}\right)-B_{p-1}\right)\\ \equiv&m^{p-2}\left(\sum_{n=1}^{m}B_{p-1}\left(\left\{\frac{pn}{m}\right\}\right)-mB_{p-1}\right)\\ \equiv&\sum_{r=0}^{m-1}m^{p-2}B_{p-1}\left(\frac{r}{m}\right)-m^{p-1}B_{p-1}=(1-m^{p-1})B_{p-1}\pmod p \end{aligned}$$

where we have applied Raabe's theorem in the last step. It is well known that $pB_{p-1}\equiv-1\pmod p$ (cf. [IR, p. 233]). Also,

$$\begin{aligned} &\sum_{n=1}^{m}\sum_{r=1}^{n}K_p(r,m)=\sum_{r=1}^{m}(m-(r-1))K_p(r,m)\\ \equiv&-\sum_{r=1}^{m}(m+1-r)K_p(m+1-r,m)=-\sum_{s=1}^{m}sK_p(s,m)\pmod p. \end{aligned}$$

So we have (2.4). $\square$

Remark. It can be shown that (2.4) is equivalent to a formula of Lerch [L] which was deduced in a different way.

3. PROOFS OF THEOREMS 1.1–1.3

Proof of Theorem 1.2. For each positive integer d we set

$$\psi(d)=\prod_{\substack{0<c<d/2\\(c,d)=1}}\binom{p-1}{\lfloor pc/d\rfloor},$$

where $\psi(1)$ and $\psi(2)$ are considered as 1. For any $a\in\mathbb{Z}$ with $p\nmid a$, clearly

$$\begin{aligned}a^p-a=&a\left(a^{(p-1)/2}+\left(\frac{a}{p}\right)\right)\left(a^{(p-1)/2}-\left(\frac{a}{p}\right)\right)\\ \equiv&2a\left(\frac{a}{p}\right)\left(a^{(p-1)/2}-\left(\frac{a}{p}\right)\right)\ (\mathrm{mod}\ p^2).\end{aligned}$$

Thus, Theorem 1.1 of [S1] implies that if $d\not\equiv0\ (\mathrm{mod}\ p)$ then

$$\begin{aligned}&(-1)^{\frac{p-1}{2}\lfloor\frac{d-1}{2}\rfloor}\prod_{0<c<d/2}\binom{p-1}{\lfloor pc/d\rfloor}\\ \equiv&\begin{cases}(\frac{d}{p})+(\frac{d}{p})\frac{d^p-d}{2}&\text{if }2\nmid d,\\ (\frac{2d}{p})+(\frac{2d}{p})\frac{d^p-d}{2}-(\frac{2d}{p})\frac{2^p-2}{2}&\text{if }2\mid d,\end{cases}\\ \equiv&\left(\frac{d}{p}\right)\left(\frac{2}{p}\right)^{[2|d]}\left(1+\frac{d^p-d}{2}-[2\mid d](2^{p-1}-1)\right)\ \ (\mathrm{mod}\ p^2).\end{aligned}$$

Since $\prod_{0<k<n/2}\binom{p-1}{\lfloor pk/n\rfloor}=\prod_{d\mid n}\psi(d)$ for $n=1,2,\ldots$, applying the Möbius inversion formula we get that

$$\begin{aligned}\psi(m)=&\prod_{d\mid m}\prod_{0<c<d/2}\binom{p-1}{\lfloor pc/d\rfloor}^{\mu(m/d)}\\ \equiv&(-1)^{\frac{p-1}{2}\sum_{d\mid m}\mu(\frac{m}{d})(\frac{d-1}{2}-\frac{[2|d]}{2})}\left(\frac{2}{p}\right)^{\sum_{d\mid m}\mu(m/d)[2|d]}\\ &\times\prod_{d\mid m}\left(\frac{d}{p}\right)^{\mu(m/d)}\times\prod_{d\mid m}\left(1+\mu\left(\frac{m}{d}\right)\left(\frac{d^p-d}{2}-[2\mid d](2^{p-1}-1)\right)\right)\ (\mathrm{mod}\ p^2).\end{aligned}$$

By elementary number theory, $\sum_{d\mid m}\mu(\frac{m}{d})\frac{d-1}{2}=\frac{\varphi(m)}{2}$ and also

$$\sum_{d\mid m}\mu\left(\frac{m}{d}\right)[2\mid d]=\sum_{2c\mid m}\mu\left(\frac{m}{2c}\right)=[2\mid m]\sum_{c\mid(m/2)}\mu\left(\frac{m/2}{c}\right)=0$$

since $m > 2$. Therefore

$$(-1)^{\frac{\varphi(m)}{2}\cdot\frac{p-1}{2}}\psi(m) \equiv \prod_{d\mid m}\left(\frac{d}{p}\right)^{\mu(m/d)} \times \left(1+\sum_{d\mid m}\mu\left(\frac{m}{d}\right)\frac{d^p-d}{2}\right) \pmod{p^2}.$$

Observe that

$$\begin{aligned}\prod_{d\mid m}\left(\frac{d}{p}\right)^{\mu(m/d)} &= \prod_{I\subseteq\{1,\dots,r\}}\left(\frac{m/\prod_{i\in I}p_i}{p}\right)^{\mu(\prod_{i\in I}p_i)}\\
=\left(\frac{m^{2^r}/\prod_{I\subseteq\{1,\dots,r\}}\prod_{i\in I}p_i}{p}\right) &= \left(\frac{m^{2^r}/\prod_{i=1}^r p_i^{2^{r-1}}}{p}\right)\\
=\left(\frac{\prod_{i=1}^r p_i^{2^{r-1}(2\alpha_i-1)}}{p}\right) &= \left(\frac{p_1\cdots p_r}{p}\right)^{2^{r-1}} = \left(\frac{p_1}{p}\right)^{[r=1]}.\end{aligned}$$

Also,

$$\begin{aligned}&\varphi(m)+\sum_{d\mid m}\mu\left(\frac{m}{d}\right)(d^p-d) = \sum_{d\mid m}\mu(d)\frac{m^p}{d^p} = m^p\prod_{i=1}^r\left(1-p_i^{-p}\right)\\
=&\prod_{i=1}^r\left(p_i^{\alpha_i p}-p_i^{(\alpha_i-1)p}\right) = \prod_{i=1}^r\left((p_i+(p_i^p-p_i))^{\alpha_i}-(p_i+(p_i^p-p_i))^{\alpha_i-1}\right)\\
\equiv&\prod_{i=1}^r\left(p_i^{\alpha_i}+\alpha_i p_i^{\alpha_i-1}(p_i^p-p_i)-\left(p_i^{\alpha_i-1}+(\alpha_i-1)p_i^{\alpha_i-2}(p_i^p-p_i)\right)\right)\\
\equiv&\prod_{i=1}^r\left(\varphi(p_i^{\alpha_i})+(p_i^{p-1}-1)(\alpha_i p_i^{\alpha_i}-(\alpha_i-1)p_i^{\alpha_i-1})\right)\\
\equiv&\varphi(m)\left(1+\sum_{i=1}^r\frac{p_i^{p-1}-1}{p_i-1}(\alpha_i p_i-\alpha_i+1)\right) \pmod{p^2}.\end{aligned}$$

Thus (1.3) holds in view of the above. □

Proof of Theorem 1.3. Write $\varepsilon^{p-(\frac{d}{p})} = (V+U\sqrt{d})/2$ where $U, V \in \mathbb{Z}$, and let p' be an integer with $pp' \equiv 1 \pmod d$. Theorem 3.1 of Williams [W] states that

$$h\frac{U}{p} \equiv -\left(\frac{d}{p}\right)N(\varepsilon)^{((\frac{d}{p})-1)/2}\sum_{i=1}^{p-1}\frac{\beta_p(i)}{i} \pmod p$$

where $\beta_p(i) = \sum_{0<j<d\{p'i/d\}}(\frac{d}{j})$.

Let $\bar\varepsilon = (a-b\sqrt{d})/2$. Then $\varepsilon+\bar\varepsilon = a$ and $\varepsilon\bar\varepsilon = N(\varepsilon)$. Clearly

$$v_n(a,N(\varepsilon))+u_n(a,N(\varepsilon))b\sqrt{d} = \varepsilon^n+\bar\varepsilon^n+\frac{\varepsilon^n-\bar\varepsilon^n}{\varepsilon-\bar\varepsilon}b\sqrt{d} = 2\varepsilon^n$$

for $n = 0, 1, \ldots$, thus $U = bu_{p-(\frac{d}{p})}(a, N(\varepsilon)) = u$ (and $V = v_{p-(\frac{d}{p})}(a, N(\varepsilon))$).

Observe that

$$\begin{aligned}\sum_{i=1}^{p-1}\frac{\beta_p(i)}{i} =&\sum_{j=1}^{d-1}\left(\frac{d}{j}\right)\sum_{\substack{0<i<p\\ d\{p'i/d\}>j}}\frac{1}{i} = \sum_{j=1}^{d-1}\left(\frac{d}{j}\right)\sum_{j<r<d}\sum_{\substack{0<i<p\\ d\mid p'i-r}}\frac{1}{i}\\ =&\sum_{j=1}^{d-1}\left(\frac{d}{j}\right)\sum_{j<r<d}\sum_{\substack{0<i<p\\ d\mid i-rp}}\frac{1}{i} = \sum_{j=1}^{d-1}\left(\frac{d}{j}\right)\sum_{j<r<d}K_p(r,d).\end{aligned}$$

As $\chi(j) = (\frac{d}{j})$ is a nontrivial multiplicative character modulo d, the sum $\sum_{j=1}^{d-1}(\frac{d}{j})$ vanishes. Therefore, with the help of Corollary 2.1, we have

$$\begin{aligned}\sum_{i=1}^{p-1}\frac{\beta_p(i)}{i} =&\sum_{j=1}^{d-1}\left(\frac{d}{j}\right)\left(\sum_{r=1}^{d}K_p(r,d) - \sum_{r=1}^{j}K_p(r,d)\right)\\ \equiv&\sum_{j=1}^{d-1}\left(\frac{d}{j}\right)\frac{1}{d}\left(0 - B_{p-1}\left(\left\{\frac{pj}{d}\right\}\right) + B_{p-1}\right)\\ \equiv&-\frac{1}{d}\left(\frac{d}{p}\right)\sum_{j=1}^{d-1}\left(\frac{d}{pj}\right)\left(B_{p-1}\left(\left\{\frac{pj}{d}\right\}\right) - B_{p-1}\right)\\ \equiv&-\frac{1}{d}\left(\frac{d}{p}\right)\sum_{c=1}^{d-1}\left(\frac{d}{c}\right)\left(B_{p-1}\left(\frac{c}{d}\right) - B_{p-1}\right) \pmod{p}.\end{aligned}$$

Combining the above we obtain (1.4).

For each $c = 1, \ldots, d-1$, we have $\chi(d-c) = \chi(-1)\chi(c) = \chi(c)$; also

$$\begin{aligned}(-1)^{\lfloor pc/d\rfloor}\binom{p-1}{\lfloor pc/d\rfloor} =& \prod_{k=1}^{\lfloor pc/d\rfloor}\left(1-\frac{p}{k}\right)\\ \equiv&1 - p\sum_{k=1}^{\lfloor pc/d\rfloor}\frac{1}{k} \equiv 1 + p\left(B_{p-1}\left(\left\{\frac{pc}{d}\right\}\right) - B_{p-1}\right) \pmod{p^2}.\end{aligned}$$

Taking the above congruence and (1.3) modulo p, we obtain

$$\begin{aligned}\prod_{\substack{0<c<d/2\\ (c,d)=1}}(-1)^{\lfloor pc/d\rfloor} \equiv& \prod_{\substack{0<c<d/2\\ (c,d)=1}}\binom{p-1}{\lfloor pc/d\rfloor}\\ \equiv&(-1)^{\frac{\varphi(d)}{2}\cdot\frac{p-1}{2}}\left(\frac{d}{p}\right)^{[d\text{ is a prime power}]} \pmod{p}\end{aligned}$$

and hence

$$\prod_{0<c<d/2}(-1)^{\lfloor pc/d\rfloor(\frac{d}{c})}=\left(\frac{d}{p}\right)^{[d=8\ \text{or}\ d\in\mathbb{P}]}.$$

(Note that $4\mid\varphi(d)$ and no square of an odd prime divides d.) On the other hand,

$$\begin{aligned}&\prod_{0<c<d/2}\left((-1)^{\lfloor pc/d\rfloor}\binom{p-1}{\lfloor pc/d\rfloor}\right)^{(\frac{d}{c})}\\ \equiv&\prod_{0<c<d/2}\left(1+p\left(\frac{d}{c}\right)\left(B_{p-1}\left(\left\{\frac{pc}{d}\right\}\right)-B_{p-1}\right)\right)\\ \equiv&1+\frac{p}{2}\sum_{0<c<d/2}\left(\frac{d}{c}\right)\left(B_{p-1}\left(\left\{\frac{pc}{d}\right\}\right)-B_{p-1}\right)\\ &+\frac{p}{2}\sum_{0<c<d/2}\left(\frac{d}{d-c}\right)\left(B_{p-1}\left(\left\{\frac{p(d-c)}{d}\right\}\right)-B_{p-1}\right)\\ \equiv&1+\frac{p}{2}\sum_{c=1}^{d-1}\left(\frac{d}{c}\right)\left(B_{p-1}\left(\left\{\frac{pc}{d}\right\}\right)-B_{p-1}\right)\\ \equiv&1+\frac{p}{2}\left(\frac{d}{p}\right)\sum_{r=1}^{d-1}\left(\frac{d}{r}\right)\left(B_{p-1}\left(\frac{r}{d}\right)-B_{p-1}\right)\pmod{p^2}.\end{aligned}$$

These, together with (1.4), yield

$$\prod_{0<c<d/2}\binom{p-1}{\lfloor pc/d\rfloor}^{(\frac{d}{c})}\equiv\left(\frac{d}{p}\right)^{[d=8\ \text{or}\ d\in\mathbb{P}]}\left(1+\frac{dhu}{2}\left(\frac{d}{p}\right)^{[N(\varepsilon)=1]}\right)\pmod{p^2}.$$

It is well known that $N(\varepsilon)=-1$ if $d=8$ or $d\in\mathbb{P}$ (see, e.g., [C, pp. 185-186]). So the desired (1.5) follows. □

Proof of Theorem 1.1. By Theorem 1.2 and the proof of Theorem 1.3,

$$\left(\frac{d}{p}\right)^{[d=8\ \text{or}\ d\in\mathbb{P}]}\prod_{\substack{0<c<d/2\\(c,d)=1}}\binom{p-1}{\lfloor pc/d\rfloor}\equiv1+\frac{\varphi(d)}{2}F(d,p)\pmod{p^2}$$

where

$$\begin{aligned}F(d,p)=&[\alpha>0](2\alpha-\alpha+1)\frac{2^{p-1}-1}{2-1}+\sum_{0<i\leqslant r}(p_i-1+1)\frac{p_i^{p-1}-1}{p_i-1}\\ =&(\alpha+[\alpha>0])(2^{p-1}-1)+\sum_{0<i\leqslant r}\frac{p_i^p-p_i}{p_i-1};\end{aligned}$$

also

$$\left(\frac{d}{p}\right)^{[d=8\text{ or }d\in\mathbb{P}]}\prod_{\substack{0<c<d/2\\(c,d)=1}}\binom{p-1}{\lfloor pc/d\rfloor}^{(\frac{d}{c})}\equiv 1+\frac{dhu}{2}\left(\frac{d}{p}\right)^{[N(\varepsilon)=1]}\pmod{p^2}$$

where $u=bu_{p-(\frac{d}{p})}(a,N(\varepsilon))\equiv 0\pmod{p}$. Therefore

$$\begin{aligned}\prod_{\substack{0<c<d/2\\(\frac{d}{c})=\rho}}\binom{p-1}{\lfloor pc/d\rfloor}\binom{p-1}{\lfloor p(d-c)/d\rfloor}&=\prod_{\substack{0<c<d/2\\(c,d)=1}}\binom{p-1}{\lfloor pc/d\rfloor}^{1+\rho(\frac{d}{c})}\\&\equiv\left(1+\frac{\varphi(d)}{2}F(d,p)\right)\left(1+\frac{dhu}{2}\left(\frac{d}{p}\right)^{[N(\varepsilon)=1]}\right)^{\rho}\\&\equiv\left(1+\frac{\varphi(d)}{2}F(d,p)\right)\left(1+\rho\frac{dhu}{2}\left(\frac{d}{p}\right)^{[N(\varepsilon)=1]}\right)\\&\equiv 1+\frac{\varphi(d)}{2}F(d,p)+\rho\frac{dhu}{2}\left(\frac{d}{p}\right)^{[N(\varepsilon)=1]}\pmod{p^2}.\end{aligned}$$

This proves (1.2). We are done. □

Acknowledgment. The author thanks the referee for his many helpful comments.

REFERENCES

[C] H. Cohn, *Advanced Number Theory*, Dover Publ. Inc., New York, 1962.

[CP] R. Crandall and C. Pomerance, *Prime Numbers: A Computational Perspective*, Springer, New York, 2001.

[DSS] K. Dilcher, L. Skula and I. Sh. Slavutskii, *Bernoulli numbers, 1713/1990*, Queen's Papers in Pure and Appl. Math. **87**(1990). The website of the on-line version is `http://www.mathstat.dal.ca/~dilcher/bernoulli.html`.

[G] A. Granville, *Arithmetic properties of binomial coefficients.I. Binomial coefficients modulo prime powers,* in: Organic mathematics (Burnaby, BC, 1995), 253–276, CMS Conf. Proc., 20, Amer. Math. Soc., Providence, RI, 1997.

[GS] A. Granville and Z. W. Sun, *Values of Bernoulli polynomials*, Pacific J. Math. **172** (1996), 117–137.

[IR] K. Ireland and M. Rosen, *A Classical Introduction to Modern Number Theory* (Graduate texts in math.; 84), 2nd ed., Springer, New York, 1990.

[L] M. Lerch, *Zur Theorie des Fermatschen Quotienten* $(a^{p-1}-1)/p=q(a)$, Math. Ann. **60** (1905), 471–490.

[R] P. Ribenboim, *The Book of Prime Number Records*, Springer, New York, 1988.

[S1] Z. W. Sun, *Products of binomial coefficients modulo* p^2, Acta Arith. **97** (2001), 87–98.

[S2] Z. W. Sun, *On the sum* $\sum_{k\equiv r\pmod{m}}\binom{n}{k}$ *and related congruences*, Israel J. Math. **128** (2002), 135–156.

[S3] Z. W. Sun, *General congruences for Bernoulli polynomials*, Discrete Math. **262** (2003), 253–276.

[W] H. C. Williams, *Some formulae concerning the fundamental unit of a real quadratic field*, Discrete Math. **92** (1991), 431–440.